

DUNFORD-PETTIS OPERATORS ON THE SPACE OF BOCHNER INTEGRABLE FUNCTIONS

MARIAN NOWAK

*Faculty of Mathematics, Computer Science and Econometrics
University of Zielona Góra
Szafrana 4A, 65-516 Zielona Góra, Poland
e-mail: M.Nowak@wmie.uz.zgora.pl*

Abstract. Let (Ω, Σ, μ) be a finite measure space and let X be a real Banach space. Let $L^\Phi(X)$ be the Orlicz-Bochner space defined by a Young function Φ . We study the relationships between Dunford-Pettis operators T from $L^1(X)$ to a Banach space Y and the compactness properties of the operators T restricted to $L^\Phi(X)$. In particular, it is shown that if X is a reflexive Banach space, then a bounded linear operator $T : L^1(X) \rightarrow Y$ is Dunford-Pettis if and only if T restricted to $L^\infty(X)$ is $(\tau(L^\infty(X), L^1(X^*)), \|\cdot\|_Y)$ -compact.

1. Introduction and preliminaries. Recall that a bounded linear operator T between two Banach spaces is a Dunford-Pettis operator if T maps weakly convergent sequences onto norm convergent sequences. J. Bourgain [B, Proposition 1] showed that a bounded linear operator T from L^1 to a Banach space Y is a Dunford-Pettis operator if and only if T restricted to L^p for some $p \in (1, \infty]$ is compact. The purpose of this paper is to extend and strengthen this result for operators defined on the space of Bochner integrable functions $L^1(X)$. We study the relationships between Dunford-Pettis operators $T : L^1(X) \rightarrow Y$ and the compactness properties of T restricted to Orlicz-Bochner spaces $L^\Phi(X)$ (see Theorem 2.1, Theorem 2.3 and Corollary 2.5 below).

We denote by $\sigma(L, K)$ the weak topology on L with respect to the dual pair $\langle L, K \rangle$. Let (L, ξ) and (M, η) be Hausdorff locally convex spaces. Recall that a linear operator $S : L \rightarrow M$ is (ξ, η) -compact if there exists a neighbourhood U of 0 for ξ such that $S(U)$ is a relatively compact set in (M, η) . By $\text{Bd}(L, \xi)$ we denote the collection of all ξ -bounded sets in L . Moreover, $(L, \xi)^*$ stands for the topological dual of (L, ξ) .

For terminology and basic properties concerning Banach function spaces we refer to [KA]. Now we recall terminology concerning Orlicz space (see [Lu], [RR] for more details). From now

2010 *Mathematics Subject Classification*: 47B38, 47B05, 46E40, 46A70.

Key words and phrases: Orlicz-Bochner spaces, mixed topologies, generalized DF-spaces, Mackey topologies, Dunford-Pettis operators, compact operators

The paper is in final form and no version of it will be published elsewhere.

we assume that (Ω, Σ, μ) is a finite measure space. By a Young function we mean here a non-zero convex, left continuous function $\Phi : [0, \infty) \rightarrow [0, \infty]$ that is vanishing and continuous at 0. We say that Φ jumps to infinity, if $\Phi(t) = \infty$ for all $t \geq t_0 > 0$.

The Orlicz space $L^\Phi = \{u \in L^0 : \int_\Omega \Phi(\lambda|u(\omega)|)d\mu < \infty \text{ for some } \lambda > 0\}$ can be equipped with the complete Riesz norm:

$$\|u\|_\Phi = \inf \left\{ \lambda > 0 : \int_\Omega \Phi(|u(\omega)|/\lambda) d\mu \leq 1 \right\}.$$

Then L^Φ is a perfect Banach function space and $L^\infty \subset L^\Phi \subset L^1$, where the inclusion maps are continuous. Moreover, the Köthe dual $(L^\Phi)'$ of L^Φ is equal to the Orlicz space L^{Φ^*} , where Φ^* stands for the Young function complementary to Φ in the sense of Young. The associated norm $\|\cdot\|_{\Phi^*}^0$ on L^{Φ^*} (called the Orlicz norm) can be defined by

$$\|\cdot\|_{\Phi^*}^0 = \sup \left\{ \int_\Omega |u(\omega)v(\omega)| d\mu : u \in L^\Phi, \|u\|_\Phi \leq 1 \right\}.$$

Note that if $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = \infty$, then $L^\Phi \subsetneq L^1$ and

$$L^\infty \subsetneq (L^{\Phi^*})_a = E^{\Phi^*} = \left\{ v \in L^{\Phi^*} : \int_\Omega \Phi(\lambda|v(\omega)|) d\mu < \infty \text{ for all } \lambda > 0 \right\}.$$

In particular, if Φ jumps to infinity, then $L^\Phi = L^\infty$. If $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} < \infty$, then $L^\Phi = L^1$ and $L^{\Phi^*} = L^\infty$.

From now on we assume that $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are real Banach spaces and X^* , Y^* denote their Banach duals. By $L^0(X)$ we denote the set of μ -equivalence classes of all strongly Σ -measurable functions $f : \Omega \rightarrow X$.

For $f \in L^0(X)$ let $\tilde{f}(\omega) = \|f(\omega)\|_X$ for $\omega \in \Omega$. Then the space

$$L^\Phi(X) = \{f \in L^0(X) : \tilde{f} \in L^\Phi\}$$

provided with the norm $\|f\|_{L^\Phi(X)} := \|\tilde{f}\|_\Phi$ is a Banach space and is usually called an *Orlicz-Bochner space* (see [CM], [L], [RR] for more details).

Now we recall terminology and basic results concerning duality of the spaces $L^\Phi(X)$ (see [Bu₁], [Bu₂]). A linear functional F on $L^\Phi(X)$ is said to be *order continuous* if $F(f_\alpha) \rightarrow 0$ whenever $\tilde{f}_\alpha \xrightarrow{(o)} 0$ in L^Φ . The set of all order continuous functionals on $L^\Phi(X)$ will be denoted by $L^\Phi(X)_n^\sim$ and called the order continuous dual of $L^\Phi(X)$. Then $L^\Phi(X)^* = L^\Phi(X)_n^\sim$ if Φ satisfies the so called Δ_2 -condition, i.e., $\limsup_{t \rightarrow \infty} \frac{\Phi(2t)}{\Phi(t)} < \infty$. Due to Bukhvalov (see [Bu₁], [Bu₂]) if X^* has the Radon-Nikodym property (in particular, X is reflexive), then $L^\Phi(X)_n^\sim$ can be identified with $L^{\Phi^*}(X^*)$ throughout the mapping: $L^{\Phi^*}(X^*) \ni g \mapsto F_g \in L^\Phi(X)_n^\sim$, where

$$F_g(f) = \int_\Omega \langle f(\omega), g(\omega) \rangle d\mu \quad \text{for all } f \in L^\Phi(X).$$

Note that $L^1(X)_n^\sim = L^1(X)^* = \{F_g : g \in L^\infty(X^*)\}$ if X is reflexive.

For a subset H of $L^\Phi(X)$ let $\tilde{H} = \{\tilde{f} : f \in H\}$. By $B_{L^\Phi(X)}$ (resp. B_{L^Φ}) we will denote closed unit ball in $(L^\Phi(X), \|\cdot\|_{L^\Phi(X)})$ (resp. $(L^\Phi, \|\cdot\|_\Phi)$). Then $\tilde{B}_{L^\Phi(X)} = B_{L^\Phi}$.

The following characterization of relative $\sigma(L^\Phi(X), L^\Phi(X)_n^\sim)$ -compactness in $L^\Phi(X)$ will be of importance (see [N₁, Theorem 2.7, Proposition 2.1]).

PROPOSITION 1.1. *Assume that X is a reflexive Banach space and Φ is a Young function. Then for a subset H of $L^\Phi(X)$ the following statements are equivalent:*

- (i) H is relatively $\sigma(L^\Phi(X), L^{\Phi^*}(X^*))$ -compact.
- (ii) \tilde{H} is relatively $\sigma(L^\Phi, L^{\Phi^*})$ -compact.

(iii) The functional $p_{\tilde{H}}$ on L^{Φ^*} defined by $p_{\tilde{H}}(v) = \sup_{u \in \tilde{H}} \int_{\Omega} |u(\omega)v(\omega)| d\mu$ is an order continuous seminorm.

2. Dunford-Pettis operators on $L^1(X)$. We study the relationships between Dunford-Pettis operators $T : L^1(X) \rightarrow Y$ and the compactness properties of the operator T restricted to $L^{\Phi}(X)$. Note that a bounded linear operator $T : L^1(X) \rightarrow Y$ is a Dunford-Pettis operator if and only if T maps relatively weakly compact sets in $L^1(X)$ onto relatively norm compact sets in Y (see [AB, § 19]).

Let $i_{\Phi} : L^{\Phi}(X) \rightarrow L^1(X)$ stand for the inclusion map.

THEOREM 2.1. *Let $T : L^1(X) \rightarrow Y$ be a bounded linear operator. Assume that Φ is Young function and let $T \circ i_{\Phi} : L^{\Phi}(X) \rightarrow Y$ be a $(\|\cdot\|_{\Phi}, \|\cdot\|_Y)$ -compact operator. Then T is a Dunford-Pettis operator.*

Proof. We see that $T(B_{L^{\Phi}(X)})$ is relatively compact in $(Y, \|\cdot\|_Y)$. Let H be a relatively $\sigma(L^1(X), L^1(X)^*)$ -compact subset of $L^1(X)$. To show that $T(H)$ is relatively compact in $(Y, \|\cdot\|_Y)$ it is enough to show in view of [D, p. 5], that for every $\varepsilon > 0$ there exists a relatively compact subset K_{ε} of $(Y, \|\cdot\|_Y)$ such that

$$T(H) \subset \varepsilon B_Y + K_{\varepsilon}.$$

where B_Y is a closed unite ball in Y . Note that the set \tilde{H} is uniformly integrable in L^1 (see [DU, Theorem 4, p. 104]). For $f \in L^1(X)$ and $\lambda > 0$ let

$$A_{f,\lambda} = \{\omega \in \Omega : \tilde{f}(\omega) > \lambda\}.$$

Then

$$\lim_{\lambda \rightarrow \infty} \sup_{f \in H} \int_{A_{f,\lambda}} \tilde{f}(\omega) d\mu = \lim_{\lambda \rightarrow \infty} \sup_{f \in H} \|1_{A_{f,\lambda}} f\|_{L^1(X)} = 0.$$

Let $\varepsilon > 0$ be given. Then there exists $\lambda_{\varepsilon} > 0$ such that for each $f \in H$ we have

$$\|1_{A_{f,\lambda_{\varepsilon}}} f\|_{L^1(X)} \leq \frac{\varepsilon}{\|T\|}.$$

Hence for $f \in H$ we get

$$\|T(1_{A_{f,\lambda_{\varepsilon}}} f)\|_Y \leq \|T\| \cdot \|1_{A_{f,\lambda_{\varepsilon}}} f\|_{L^1(X)} \leq \varepsilon.$$

Moreover, $1_{\Omega \setminus A_{f,\lambda_{\varepsilon}}}(\omega) \tilde{f}(\omega) \leq \lambda_{\varepsilon}$ for $\omega \in \Omega$, so $1_{\Omega \setminus A_{f,\lambda_{\varepsilon}}} f \in L^{\infty}(X) \subset L^{\Phi}(X)$. Since $\|h\|_{L^{\Phi}(X)} \leq a \|h\|_{L^{\infty}(X)}$ for some $a > 0$ and all $h \in L^{\infty}(X)$, we get

$$\|1_{\Omega \setminus A_{f,\lambda_{\varepsilon}}} f\|_{L^{\Phi}(X)} \leq a \lambda_{\varepsilon}.$$

Hence

$$T(f) = T(1_{A_{f,\lambda_{\varepsilon}}} f) + T(1_{\Omega \setminus A_{f,\lambda_{\varepsilon}}} f) \in \varepsilon B_Y + a \lambda_{\varepsilon} T(B_{L^{\Phi}(X)}).$$

This means that the set $T(H)$ is relatively compact in $(Y, \|\cdot\|_Y)$, as desired. ■

From now we assume that Φ is a Young function such that $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = \infty$. Let \mathcal{T}_{Φ} be the topology on $L^{\Phi}(X)$ generated by the norm $\|\cdot\|_{L^{\Phi}(X)}$ on $L^{\Phi}(X)$, and let \mathcal{T}_0 stand for the complete F -norm $\|\cdot\|_{L^0(X)}$ -topology on $L^0(X)$ that generates convergence in measure. Then the mixed topology $\gamma[\mathcal{T}_{\Phi}, \mathcal{T}_0]_{L^{\Phi}(X)}$ (briefly, γ_{Φ}) on $L^{\Phi}(X)$ is the finest Hausdorff locally convex topology on $L^{\Phi}(X)$ which agrees with $\mathcal{T}_0|_{L^{\Phi}(X)}$ on $\|\cdot\|_{L^{\Phi}(X)}$ -bounded subsets of $L^{\Phi}(X)$ (see [W, 2.2.2], [F₁, Theorem 3.3]). Moreover, we have (see [F₂, Proposition 2.1]):

$$(2.1) \quad \text{Bd}(L^{\Phi}(X), \gamma_{\Phi}) = \text{Bd}(L^{\Phi}(X), \|\cdot\|_{L^{\Phi}(X)}).$$

This means that $(L^\Phi(X), \gamma_\Phi)$ is a generalized DF-space (see [Ru, Definition 1.1]).

It is known that a linear operator $T : L^\Phi(X) \rightarrow Y$ is $(\gamma_\Phi, \|\cdot\|_Y)$ -continuous if and only if T is $(\gamma_\Phi, \|\cdot\|_Y)$ -linear, i.e., $\|T(f_n)\|_Y \rightarrow 0$ whenever $\|f_n\|_{L^0(X)} \rightarrow 0$ and $\sup_n \|f_n\|_{L^\Phi(X)} < \infty$ (see [W, Theorem 2.6.1 (iii)], [F₂, Proposition 2.3]).

We shall need the following lemma.

LEMMA 2.2. *Assume that Φ is a Young function such that $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = \infty$ and X is a reflexive Banach space. Then $i_\Phi : L^\Phi(X) \rightarrow L^1(X)$ is a $(\|\cdot\|_{L^\Phi(X)}, \sigma(L^1(X), L^1(X)^*))$ -compact operator.*

Proof. To show that $B_{L^\Phi(X)}$ is a relatively $\sigma(L^1(X), L^1(X)^*)$ -compact subset of $L^1(X)$, in view of Proposition 1.1 it is enough to show that B_{L^Φ} is relatively $\sigma(L^1, L^\infty)$ -compact in L^1 , that is, the seminorm on L^∞ defined by

$$p_{B_{L^\Phi}}(v) := \sup_{u \in B_{L^\Phi}} \int_{\Omega} |u(\omega)v(\omega)| d\mu$$

is order continuous. Indeed, note that $p_{B_{L^\Phi}}(v) = \|\cdot\|_{\Phi^*}^0$ for $v \in L^\infty$, where $L^\infty \subsetneq E^{\Phi^*} = (L^{\Phi^*})_a$. Thus the proof is complete. ■

Now we are ready to proof our main result.

THEOREM 2.3. *Assume that Φ is a Young function such that $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = \infty$ and X is a reflexive Banach space. Let $T : L^1(X) \rightarrow Y$ be a Dunford-Pettis operator. Then the operator $T \circ i_\Phi : L^\Phi(X) \rightarrow Y$ is $(\gamma_\Phi, \|\cdot\|_Y)$ -compact.*

Proof. Since X is supposed to be reflexive, in view of [F₁, Theorem 3.2] we have

$$(L^\Phi(X), \gamma_\Phi)^* = \{F_g : g \in E^{\Phi^*}(X^*)\}.$$

First, we shall show that $T \circ i_\Phi : L^\Phi(X) \rightarrow Y$ is $(\gamma_\Phi, \|\cdot\|_Y)$ -linear. Indeed, let (f_n) be a sequence in $L^\Phi(X)$ such that $\|f_n\|_{L^0(X)} \rightarrow 0$ and $\sup_n \|f_n\|_{L^\Phi(X)} < \infty$. Then $f_n \rightarrow 0$ for γ_Φ (see [F₁, Theorem 3.1]), and it follows that $f_n \rightarrow 0$ for $\sigma(L^\Phi(X), E^{\Phi^*}(X^*))$ because $\sigma(L^\Phi(X), E^{\Phi^*}(X^*)) \subset \gamma_\Phi$. Hence $f_n \rightarrow 0$ for $\sigma(L^1(X), L^1(X)^*)$ because $\sigma(L^1(X), L^1(X)^*) = \sigma(L^1(X), L^\infty(X^*))$ and $L^\Phi(X) \subset L^1(X)$ and $L^\infty(X^*) \subset E^{\Phi^*}(X^*)$. Since T is a Dunford-Pettis operator, we get $\|T(f_n)\|_Y \rightarrow 0$. This means that $T \circ i_\Phi$ is $(\gamma_{L^\Phi(X)}, \|\cdot\|_Y)$ -continuous.

By Lemma 2.2 the mapping $T \circ i_\Phi$ is $(\|\cdot\|_{L^\Phi(X)}, \|\cdot\|_Y)$ -compact. Hence, in view of (2.1) $T \circ i_\Phi$ transforms γ_Φ -bounded sets in $L^\Phi(X)$ onto relatively $\|\cdot\|_Y$ -compact sets in Y . Making use of [Ru, Theorem 3.1] we conclude that $T \circ i_\Phi$ is $(\gamma_\Phi, \|\cdot\|_Y)$ -compact, as desired. ■

As an application of Theorem 2.1 and Theorem 2.3 we get:

COROLLARY 2.4. *Assume that Φ is a Young function such that $\lim_{t \rightarrow \infty} \frac{\Phi(f)}{t} = \infty$ and X is a reflexive Banach space. Then for a bounded linear operator $T : L^1(X) \rightarrow Y$ the following statements are equivalent:*

- (i) T is a Dunford-Pettis operator.
- (ii) $T \circ i_\Phi : L^\Phi(X) \rightarrow Y$ is $(\gamma_\Phi, \|\cdot\|_Y)$ -compact.
- (iii) $T \circ i_\Phi : L^\Phi(X) \rightarrow Y$ is $(\|\cdot\|_{L^\Phi(X)}, \|\cdot\|_Y)$ -compact.

In particular, if X is reflexive, then the mixed topology γ_∞ on $L^\infty(X)$ coincides with the Mackey topology $\tau(L^\infty(X), L^1(X^*))$ (see [N₂, Corollary 4.4]). Hence, as a consequence of Corollary 2.4 we get:

COROLLARY 2.5. *Assume that X is a reflexive Banach space. Then for a bounded linear operator $T : L^1(X) \rightarrow Y$ the following statements are equivalent:*

- (i) *T is a Dunford-Pettis operator.*
- (ii) *$T \circ i_\infty : L^\infty(X) \rightarrow Y$ is $(\tau(L^\infty(X), L^1(X^*)), \|\cdot\|_Y)$ -compact.*
- (iii) *$T \circ i_\infty : L^\infty(X) \rightarrow Y$ is $(\|\cdot\|_{L^\infty(X)}, \|\cdot\|_Y)$ -compact.*

References

- [AB] C.D. Aliprantis and O. Burkinshaw, *Positive Operators*, Academic Press, New York, 1985.
- [B] J. Bourgain, Dunford-Pettis operators on L^1 and the Radon-Nikodym property, *Israel J. Math.*, 37, no. 1–2 (1980), 34–47.
- [Bu₁] A.V. Bukhvalov, The analytic representation of linear operators with abstract norm (Russian), *Izv. Vyss. Vceb. Zaved. Matematika*, 11 (1975), 21–32.
- [Bu₂] A.V. Bukhvalov, The analytic representation of linear operators by means of measurable vector-valued functions (Russian), *Izv. Vyss. Vceb. Zaved. Matematika*, 7 (1977), 21–31.
- [CM] P. Cembranos and J. Mendoza, *Banach spaces of vector-valued functions*, *Lectures Notes in Math.*, 1676, Springer Verlag, Berlin, Heidelberg, 1997.
- [D] J. Diestel, *Sequences and Series in Banach spaces*, *Graduate Texts in Math.*, Springer-Verlag, New York, 1984.
- [DU] J. Diestel and J.J. Uhl, *Vector Measures*, *Amer. Math. Soc., Math. Surveys*, no. 15, Providence, Rhode Island 1977.
- [F₁] K. Feledziak, Uniformly Lebesgue topologies on Köthe-Bochner spaces, *Comment. Math. Prace Mat.*, 37 (1997), 81–98.
- [F₂] K. Feledziak, Weakly compact operators on Köthe-Bochner spaces, *Banach Center Publ.*, Vol. 79, *Proceedings of Function Spaces VIII*, Warszawa 2008, 71–77.
- [KA] L.V. Kantorovitch and A.V. Akilov, *Functional Analysis*, Pergamon Press, Oxford-Elmsford, N.Y., 1982.
- [L] Pei-Kee, Lin, *Köthe-Bochner Function Spaces*, Birkhäuser Verlag, Boston-Basel-Berlin 2003.
- [Lu] W. Luxemburg, *Banach Function Spaces*, Thesis, Assen, 1955.
- [N₁] M. Nowak, Weak sequential compactness and completeness in Köthe-Bochner spaces, *Bull. Polish Acad. Sci. Math.*, 47, no. 3 (1999), 209–220.
- [N₂] M. Nowak, Lebesgue topologies on vector-valued function spaces, *Math. Japonica*, 52, no. 2 (2000), 171–182.
- [RR] M.M. Rao and Z.D. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, New York, 1991.
- [Ru] W. Ruess, [Weakly] compact operators and DF-spaces, *Pacific J. Math.*, 98, no. 2 (1982), 419–441.
- [W] A. Wiweger, Linear spaces with mixed topology, *Studia Math.*, 20 (1961), 47–68.